WHAT IS CLAIMED IS:

Polymer 1

1. A composition of matter comprising a polymer of the general structure:

CH₃O
OCH₃
OCH₃

C/R2/D A/R1/B E/R3/F

-(CH₂)_x-/-O(CH₂)_x -/-O(CH₂)_xO-

wherein

the R1 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups;

the R2 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups;

the R3 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups;

wherein bonds A and B may independently be either ortho, meta or para with respect to the pyridyl nitrogen;

wherein bonds C and D may be either ortho, meta or para with respect one another; and

wherein bonds E and F may be either ortho, meta or para with respect one another;

wherein Y may be a moiety selected from the group consisting of $-(CH_2)_x$ - $-(CH_2)_x$ O-, $-O(CH_2)_x$ — and $-O(CH_2)_x$ O- wherein x is an integer in the range of 1 to 15 inclusive; and

wherein n is an integer greater than 1.

30

35

5

- 2. A composition according to claim 1 wherein at least one R2 substituent is a methoxy group.
- 5 3. A composition according to claim 1 wherein at least two R2 substituents are methoxy groups.
 - 4. A composition according to claim 1 wherein at least one R3 substituent is a methoxy group.
 - 5. A composition according to claim 1 wherein at least two R3 substituents are methoxy groups.
- 6. A composition according to claim 1 wherein vinyl linkage A attaches at a position ortho to the pyridyl nitrogen.
 - 7. A composition according to claim 1 wherein vinyl linkage B attaches at a position ortho to the pyridyl nitrogen.
 - 8. A composition according to claim 1 wherein vinyl linkage A attaches at a position para to the pyridyl nitrogen.
 - 9. A composition according to claim 1 wherein vinyl linkage B attaches at a position para to the pyridyl nitrogen.
 - 10. A composition according to claim 1 wherein x is an integer in the range of 1 to 6 inclusive.

Polymer 2

11. A composition of matter comprising a polymer of the general structure:

10

₩ 30 -(CH₂)_x-/-O(CH₂)_x -/-O(CH₂)_xO-

wherein

5

H C C C C C C C C C C C C C

25 □

Ш

□30 □

- the R1 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups;
- the R2 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups;
 - the R3 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups;
- the R4 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups;
 - wherein bonds A and B may be either ortho or para with respect to the pyridyl nitrogen;
 - wherein bonds C and D may independently be either ortho, meta or para with respect one another;
 - wherein bonds E and F may be either ortho, meta or para with respect one another;
 - wherein bonds G and H may be either ortho, meta or para with respect one another;
 - wherein Y may be a moiety selected from the group consisting of $-(CH_2)_x$, $-(CH_2)_x$ O-, $-O(CH_2)_x$ and $-O(CH_2)_x$ O- wherein x is an integer in the range of 1 to 15 inclusive;
 - wherein Z may be a moiety selected from the group consisting of $-(CH_2)_x$ -, $-(CH_2)_x$ O-, $-O(CH_2)_x$ and $-O(CH_2)_x$ O- wherein x is an integer in the range of 1 to 15 inclusive; and
- wherein n is an integer greater than 1.
 - 12. A composition according to claim 11 wherein at least one R2 substituent is a methoxy group.
- 40 13. A composition according to claim 11 wherein at least two R2 substituents are methoxy groups.
 - 14. A composition according to claim 11 wherein at least one R3 substituent is a methoxy group.

45

- 15. A composition according to claim 11 wherein at least two R3 substituents are methoxy groups.
- 16. A composition according to claim 11 wherein at least one R3 substituent is a methoxy group.
 - 17. A composition according to claim 11 wherein at least two R3 substituents are methoxy groups.
- 10 18. A composition according to claim 11 wherein vinyl linkage A attaches at a position ortho to the pyridyl nitrogen.
 - 19. A composition according to claim 11 wherein vinyl linkage B attaches at a position ortho to the pyridyl nitrogen.
 - 20. A composition according to claim 11 wherein vinyl linkage A attaches at a position para to the pyridyl nitrogen.
 - 21. A composition according to claim 11 wherein vinyl linkage B attaches at a position para to the pyridyl nitrogen.
 - 22. A composition according to claim 11 wherein x is an integer in the range of 1 to 6 inclusive.


Oligomers 1, 2, 3 & 4

5

15

₩ 30 23. A composition of matter comprising an oligomer of the general structure:

35 wherein



- the R2 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups;
 - the R3 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups;
- the R4 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups;
 - the R5 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups;
 - wherein bonds A and B may independently be either ortho, meta or para from one another;
 - wherein bond C may be either ortho, meta or para with respect to the respective quincyl nitrogen; and

wherein bond D may be either ortho, meta or para with respect to the respective quinoyl nitrogen.

- 24. A composition according to claim 23 wherein at least one R1 substituent is a methoxy group.
- 25. A composition according to claim 23 wherein at least two R1 substituents are methoxy groups.

Oligomer 5, 6, 7 & 8

26. A composition of matter comprising an oligomer of the general structure:

35

5

15

H

□ □20 □ 「 「 「 「 「 「 「 「 「 「 「 「 「 「 「 「 「 」 「 「 「 「 「 」 「 「 「 「 」 「 「 「 「 「 」 「 「 「 「 「 「 「 「 「 「 「 」 「 「 —

T₂₅

O

□30 N

R2 C A R1 B D R3

wherein

ļå

m

□ 30

N

35


25

□ □ 20

- the R1 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups;
- the R2 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups;
 - the R3 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups;
- wherein bonds A and B may independently be either ortho, meta or para from one another;
 - wherein bond C may be either ortho, meta or para with respect to the respective pyridyl nitrogen; and
 - wherein bond D may be either ortho, meta or para with respect to the respective pyridyl nitrogen.
 - 27. A composition according to claim 26 wherein at least one R1 substituent is a methoxy group.
 - 28. A composition according to claim 26 wherein at least two R1 substituents are methoxy groups.
 - 29. A composition according to claim 26 wherein at least one R2 substituent is a methyl group.
 - 30. A composition according to claim 26 wherein at least one R3 substituent is a methyl group.

Oligomers 9 & 10

31. A composition of matter comprising an oligomer of the general structure:



R3/R2 C A R1 B D R4/R5

5 wherein

10

ā

Ū₂₅

30

35

□20 N the R1 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups;

the R2 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups;

the R3 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups;

the R4 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups;

the R5 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups;

wherein bonds A and B may be either ortho, meta or para from one another.

- 32. A composition according to claim 31 wherein at least one R1 substituent is a methoxy group.
- 33. A composition according to claim 31 wherein at least two R1 substituents are methoxy groups.

Oligomers 11, 12 & 13

34. A composition of matter comprising an oligomer of the general structure:

wherein

10

(T

20 D N N D D 25

ΠJ

30

20

5 the R1 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups;

the R2 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups;


the R3 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups; and

wherein bonds A and B may be either ortho or para from the pyridyl nitrogen.

- A composition according to claim 34 wherein at least one R2 substituent is a 35. methoxy group.
- A composition according to claim 34 wherein two R2 substituents are methoxy groups.
- 37. A composition according to claim 34 wherein three R2 substituents are methoxy groups.
- A composition according to claim 34 wherein at least one R3 substituent is a 38. methoxy group.
 - 39. A composition according to claim 34 wherein two R3 substituents are methoxy groups.
 - 40. A composition according to claim 34 wherein three R2 substituents are methoxy groups.

Block Co-polymer of Oligomers 1, 2, 3 & 4 (Y only)

35 41. A composition of matter comprising a block co-polymer of the general structure:



wherein

5

⊨10

25

the R1 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups;

the R2 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups;

the R3 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups;

the R4 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups;

the R5 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups;

wherein bonds A and B may independently be either ortho, meta or para from one another;

wherein bond C may be either ortho, meta or para with respect to the respective quinoyl nitrogen; and

wherein bond D may be either ortho, meta or para with respect to the respective quinoyl nitrogen;

wherein Y may be a moiety attached at any point on rings R2 and R3, and may be selected from the group consisting of $-(CH_2)_x$ -, $-(CH_2)_x$ O-, $-O(CH_2)_x$ – and - $O(CH_2)_x$ O- wherein x is an integer in the range of 1 to 15 inclusive; and

wherein n is an integer greater than 1.

35

42. A composition according to claim 41 wherein at least one R1 substituent is a methoxy group.

10

HOOWHWOO.CO

₩20

25

30

Ö

ΠJ

Block Co-polymer of Oligomers 1, 2, 3 & 4 (Y, R & Z)

A composition of matter comprising a block co-polymer of the general structure:

wherein

the R1 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups;

the R2 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups;

the R3 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups;

the R4 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups:

the R5 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups;

35 the R6 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups;

wherein bonds A and B may independently be either ortho, meta or para from one another;

35

10

15

wherein bond C may be either ortho, meta or para with respect to the respective quinoyl nitrogen; and

wherein bond D may be either ortho, meta or para with respect to the respective quinoyl nitrogen;

wherein Y may be a moiety attached at any point on ring R6, and may be selected from the group consisting of $-(CH_2)_x-,-(CH_2)_xO-,-O(CH_2)_x-$ and $-O(CH_2)_xO-$ wherein x is an integer in the range of 1 to 15 inclusive;

wherein Z may be a moiety bridging any two points on rings R2 or R3 and R6, and may be selected from the group consisting of - $(CH_2)_{x^-}$,- $(CH_2)_{x}O$ -, - $O(CH_2)_{x}$ – and - $O(CH_2)_{x}O$ - wherein x is an integer in the range of 1 to 15 inclusive; and

wherein n is an integer greater than 1.

R2

C

- 45. A composition according to claim 44 wherein at least one R1 substituent is a methoxy group.
- 46. A composition according to claim 44 wherein at least two R1 substituents are methoxy groups.

Block Co-polymer of Oligomers 5, 6, 7 & 8 (Y only)

47. A composition of matter comprising a block co-polymer of the general structure:

R₁

DR3

wherein

the R1 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups;

15

the R2 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups;

- the R3 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups;
 - wherein bonds A and B may independently be either ortho, meta or para from one another;
- wherein bond C may be either ortho, meta or para with respect to the respective pyridyl nitrogen;
 - wherein bond D may be either ortho, meta or para with respect to the respective pyridyl nitrogen;
 - wherein Y may be a moiety attached at any point on ring R2, and may be selected from the group consisting of $-(CH_2)_x-,-(CH_2)_xO-, -O(CH_2)_x-$ and $-O(CH_2)_xO-$ wherein x is an integer in the range of 1 to 15 inclusive;

and

wherein n is an integer greater than 1.


- 48. A composition according to claim 47 wherein at least one R1 substituent is a methoxy group.
 - 49. A composition according to claim 47 wherein at least two R1 substituents are methoxy groups.
- 50. A composition according to claim 47 wherein at least one R2 substituent is a methyl group.
- 51. A composition according to claim 47 wherein at least one R3 substituent is a methyl group.

Block Co-polymer of Oligomers 5, 6, 7 & 8 (Y, R & Z)

52. A composition of matter comprising a block co-polymer of the general structure:

30

35



wherein 5

the R1 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups;

the R2 substituents are independently selected from the group consisting of 10 hydrogen, alkyl groups, alkoxy groups, and aryl groups;

the R3 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl-groups;

the R4 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups;

wherein bonds A and B may independently be either ortho, meta or para from one another:

wherein bond C may be either ortho, meta or para with respect to the respective pyridyl nitrogen;

25 wherein bond D may be either ortho, meta or para with respect to the respective pyridyl nitrogen;

wherein Y may be a moiety attached at any point on ring R4, and may be selected from the group consisting of $-(CH_2)_x$, $-(CH_2)_x$ O-, $-O(CH_2)_x$ – and $-O(CH_2)_x$ O- wherein x is an integer in the range of 1 to 15 inclusive;

wherein Z may be a moiety bridging any two points on rings R2 and R4, and may be selected from the group consisting of $-(CH_2)_{x-}$, $-(CH_2)_{x}O_{-}$, $-O(CH_2)_{x}$ – and -O(CH₂)_xO- wherein x is an integer in the range of 1 to 15 inclusive; and

and


wherein n is an integer greater than 1.

40 53. A composition according to claim 52 wherein at least one R1 substituent is a methoxy group.

- 54. A composition according to claim 52 wherein at least two R1 substituents are methoxy groups.
- 5 55. A composition according to claim 52 wherein at least one R2 substituent is a methyl group.
 - 56. A composition according to claim 52 wherein at least one R3 substituent is a methyl group.

Block Co-polymer of Oligomers 9 & 10 (Y only)

57. A composition of matter comprising a block co-polymer of the general structure:



wherein

the R1 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups;

the R2 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups;

the R3 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups;

the R4 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups;

the R5 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups;

10

10054565

ĪŪ20

wherein bonds A and B may be either ortho, meta or para from one another;


wherein Y may be a moiety attached at any point on rings R2 or R3, and may be selected from the group consisting of $-(CH_2)_{x}$, $-(CH_2)_{x}$ O-, $-O(CH_2)_{x}$ – and - $O(CH_2)_{x}$ O- wherein x is an integer in the range of 1 to 15 inclusive;

and

- wherein n is an integer greater than 1.
 - 58. A composition according to claim 57 wherein at least one R1 substituent is a methoxy group.
- 15 59. A composition according to claim 57 wherein at least two R1 substituents are methoxy groups.

Block Co-polymer of Oligomers 9 & 10 (Y, R and Z only)

60. A composition of matter comprising a block co-polymer of the general structure:



R3/R2 C A R1 B D R4/R5

wherein

25


30

the R1 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups;

the R2 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups;

the R3 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups;

15



- the R5 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups;
 - the R6 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups;
 - wherein bonds A and B may be either ortho, meta or para from one another;
 - wherein Y may be a moiety attached at any point on ring R6, and may be selected from the group consisting of $-(CH_2)_x-,-(CH_2)_xO-,-O(CH_2)_x-$ and $-O(CH_2)_xO-$ wherein x is an integer in the range of 1 to 15 inclusive;
 - wherein Z may be a moiety bridging any two points on rings R2 or R3 and R6, and may be selected from the group consisting of $-(CH_2)_x-,-(CH_2)_xO-,-O(CH_2)_x-$ and $-O(CH_2)_xO-$ wherein x is an integer in the range of 1 to 15 inclusive; and

and

wherein n is an integer greater than 1.